



Antioxidant and Antimicrobial Activities of Essential Oil of Lemon (*Citrus limon*) Peel *in Vitro* and in a Food Model

M.H. Moosavy^{1*}, P. Hassanzadeh¹, E. Mohammadzadeh¹, R. Mahmoudi¹, S.A. Khatibi², K. Mardani³

1. Department of Food Hygiene and Aquatic, Faculty of Veterinary Medicine, University of Tabriz, Tabriz, Iran

2. Department of Food Hygiene, Faculty of Veterinary Medicine, University of Tehran, Tehran, Iran

3. Department of Food Hygiene and Quality Control, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

HIGHLIGHTS

- The total phenol content was 81.82 ± 8.02 mg gallic acid equivalent/g of lemon peel EO.
- The total amount of flavonoids in the lemon peel EO was 11.72 ± 1.82 mg/g rutin equivalent.
- The MIC and MBC value of lemon peel EO against *Staphylococcus aureus* was 1.25 and 5%.
- Lemon peel EO showed considerable antioxidant and antimicrobial properties both *in vitro* and in food model.

Article type

Original article

Keywords

Citrus
Oils, Volatile
Antioxidants
Staphylococcus aureus

Article history

Received: 3 Nov 2016
Revised: 20 Jan 2017
Accepted: 2 March 2017

Acronyms and abbreviations

EO=Essential Oil
MIC=Minimum Inhibitory Concentration
MBC=Minimum Bactericidal Concentration
DPPH=2, 2-diphenyl-1-picrylhydrazyl
CFU=Colony Forming Unit

ABSTRACT

Background: Citrus fruits have some antioxidant and antimicrobial properties. The aim of this study was to determine the chemical compounds, antioxidant, and antimicrobial activities of Essential Oil (EO) of lemon (*Citrus limon*) peel *in vitro* and in a food model.

Methods: The analysis of the lemon peel EO was carried out using gas chromatography-mass spectrometry. Total phenolic and flavonoid content was determined using standard protocols. The antioxidant activity of the EO was also evaluated using 2, 2-diphenyl-1-picrylhydrazyl (DPPH) radical scavenging assay. Antimicrobial properties of the EO were assessed against *Staphylococcus aureus* using the broth microdilution method. Also, barley soup was chosen as food model. Data were analyzed using SPSS Inc. software version 22.0.

Results: The total phenol content was 81.82 ± 8.02 mg gallic acid equivalent/g of the EO. Also, the total amount of flavonoids in the EO of lemon peel was 11.72 ± 1.82 mg/g rutin equivalent. Lemon peel EO showed 55.09% inhibition of DPPH, showing significant difference with control group ($p < 0.05$). The MIC and MBC value of EO against *S. aureus* was 1.25 and 5%, respectively having significant difference ($p < 0.05$) with control group. A dose-dependent manner was seen in food model revealed significantly lower ($p < 0.05$) bacterial number in EO containing barley soup groups than the control one.

Conclusion: The EO of lemon peel showed considerable antioxidant and antimicrobial properties both *in vitro* and barley soup as food model.